

421-2274-1-CE

by Rahmat Catur Wibowo

General metrics

29,150

characters

4,553


words

214

sentences

18 min 12 secreading
time**35 min 1 sec**speaking
time

Score



This text scores better than 84%
of all texts checked by Grammarly

Writing Issues

171

Issues left

18

Critical

153

Advanced

Plagiarism

This text hasn't been checked for plagiarism

Writing Issues

106

Clarity

28

Wordy sentences



47

Passive voice misuse



2

Hard-to-read text



11

Paragraph can be perfected



15

Unclear sentences



3

Intricate text



55

Correctness

7

Ungrammatical sentence



2

Misspelled words



17

Punctuation in compound/complex sentences



7

Incorrect phrasing



2

Text inconsistencies



2

Comma misuse within clauses



1

Misplaced words or phrases



1

Determiner use (a/an/the/this, etc.)



8

Incorrect citation format




2

Wrong or missing prepositions



3

Improper formatting



3

Unknown words



9

Engagement

9

Word choice



1**Delivery****1**

Inappropriate colloquialisms



Unique Words**17%**

Measures vocabulary diversity by calculating the percentage of words used only once in your document

unique words

Rare Words**37%**

Measures depth of vocabulary by identifying words that are not among the 5,000 most common English words.

rare words

Word Length**3.7**

Measures average word length

characters per word

Sentence Length**21.3**

Measures average sentence length

words per sentence

421-2274-1-CE

JGE (Jurnal Geofisika Eksplorasi) XX (XXXX) XXXXXX Author et al

JGE (Jurnal Geofisika Eksplorasi) XX (XXXX) XXXXXX Author et al

3

11

JGE (Jurnal Geofisika Eksplorasi)

Vol. xx No. xx, xxxxxx xxx (xxx-xxx) <https://doi.org/XXXX/XXXX>

12

2D INVERSION MODELING OF OIL AND GAS BASINS USING GRAVITY DATA IN
TAMBELANG DISTRICT

PEMODELAN INVERSI 2D CEKUNGAN MINYAK DAN GAS MENGGUNAKAN DATA
GRAVITASI DI KECAMATAN TAMBELANG

Belista Gunawan¹, Nanda Ridki Permana^{2*}

¹ GeoXplore Indonesia, Kincir Air Street, Pondok Manggis Block B6, Bojong
Baru, Bojonggede, Bogor, West Java, Indonesia, 16920; (+62) 895337066366

² PT Minelog Services Indonesia; Bumi Serpong Damai, Techno Park Industrial &
Warehouse Zone Block G1 No. 10, Sector 11 Street, Setu Subdistrict, South
Tangerang City, Banten, Indonesia, 15220; (021) 75882193

Received: 2024, March 08th Accepted: 2024, July 23rd

Keywords:

Drilling Point;

Gravity;

Inversion;

Oil and Gas;

Reservoir.

Correspondent Email:

nandaridki836@gmail.com

How to cite this article:

Doe, J. (2021). Judul Artikel. Jurnal Geofisika Eksplorasi, 6(2), 156-168.

© 2021 JGE (Jurnal Geofisika Eksplorasi). This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY NC)

Abstract. Pertamina Hulu Energi discovered a new source of oil and gas reserves in Tambelang, Bekasi Regency, which was¹ named East Pondok Aren (EPN-001). These oil and gas reserves have great potential based on the oil flow test in the second Drill Stem Test. The drill data needs to² be correlated with other supporting data such as³ geophysical methods to ensure the accuracy³. Therefore, this research was conducted to examine⁴ the subsurface structure of oil and gas reserves in EPN-001 through 2D inversion modeling using the gravity method. Gravity satellite data was taken⁶ via the GGMPPlus website with a total of 23,898 data in the form of gravity disturbance, geoid, and topography, so only terrain and bouguer⁷ corrections were carried out⁸ to get the Complete Bouguer Anomaly value. The research results show low anomaly values from

-0.4463 – (-0.0339) mGal ⁹which ¹⁰are thought to be associated with alluvium lithology and the Lower Cibulakan Formation. In the 2D model, it can ¹¹be seen that the drilling point in the Lower Cibulakan Formation layer has a density contrast value of -0.002 – 0.001 g/cc at a depth of 2590 m, and is known to be a reservoir that stores hydrocarbon reserves in the research area.

Abstrak. Pertamina Hulu Energi menemukan sumber cadangan minyak dan gas baru di Tambelang, Kabupaten Bekasi, yang diberi nama East Pondok Aren (EPN-001). Cadangan minyak dan gas bumi tersebut sangat berpotensi berdasarkan uji aliran minyak bumi pada Drill Stem Test kedua. Data bor tersebut perlu dikorelasikan dengan data pendukung lainnya seperti metode geofisika untuk dapat memastikan keakuratannya. Oleh karena itu, penelitian ini dilakukan untuk melihat struktur bawah permukaan cadangan minyak dan gas di EPN-001 melalui pemodelan inversi 2D menggunakan metode gravitasi. Data satelit gravitasi diambil melalui website GGMPlus dengan jumlah sebanyak 23.898 data berupa gravity disturbance, geoid dan topografi, sehingga hanya dilakukan koreksi terrain dan bouguer untuk mendapat nilai Anomali Bouguer Lengkap. Hasil penelitian menunjukkan nilai anomali rendah berkisar -0.4463 – (-0.0339) mGal diduga berasosiasi dengan litologi alluvium dan formasi Cibulakan Bawah. Pada model 2D, terlihat bahwa drilling point yang terdapat di lapisan Formasi Cibulakan Bawah memiliki nilai kontras densitas -0.002 – 0.001 g/cc pada kedalaman 2590 m, dan diketahui sebagai reservoar yang menyimpan cadangan hidrokarbon di daerah penelitian.

1. INTRODUCTION

Petroleum reserves in 2019 decreased by 49.75%, while natural gas reserves decreased by 42.98% compared to 2018 (Directorate General of Oil and Gas, 2021). Based on data from the Ministry of Energy and Mineral Resources (ESDM), oil reserves in Indonesia are only available for 9.5 years and natural gas for 19.9 years, assuming no new reserves are discovered (Rahmayanti, 2021). As a result, continuous oil production and oil reserves will increasingly run out, causing the oil exploration process to stop,¹² resulting in old wells (Rosid, 2020). Meanwhile, it is important¹³ to continue oil and gas exploration in an effort to¹⁴ increase and maintain hydrocarbon production rates in Indonesia (Prasetiyohadi, 2022).

Therefore, steps are needed to increase the amount of energy reserves. One way is to optimize existing reserve potential through exploration and drilling activities in oil and gas fields, both those that have been operated¹⁵ previously and the discovery of new sources that have not been exploited¹⁶ (Dewi, 2020); (Purnama, 2020). However, in reality, the average oil and gas production achievement in the last two years has decreased due to a decline in natural reservoir performance and also¹⁷ the absence of new reserves that would replace reserves that continue to be produced (Directorate General of Oil and Gas, 2019).

PT Pertamina Hulu Energi (PHE)¹⁸ discovered a new source of oil and gas reserves in the Tambun Field WKP, Tambelang District, Bekasi Regency, West Java. The new reserve discovered is called East Pondok Aren (EPN-001) which¹⁹ is proof of a new concept in the form of a stratigraphic trap in the Lower Cibulakan Formation, Ciputat Sub Basin,²⁰ and is included²¹ in the onshore North West Java Basin (CNN Indonesia Team, 2023).

Drilling of the East Pondok Aren (EPN)-001 exploration well was started on August ²²18 2023. As a result, EPN-001 succeeded in showing that the flow of oil in the second Drill Stem Test (DST) was 402 Barrels of Oil Per Day (BOPD) and gas reached ²³1, 00 million standard cubic feet per day or Million Standard Cubic Feet per Day (MMSCFD) at a depth of 2,590 meters measured depth (mMD) (Ira, 2023).

To utilize the energy potential of oil and gas,²⁴ accurate data analysis is needed by optimizing the exploration stages in more depth. Apart from the drilled well data that has been obtained,²⁵ other supporting data is also needed to see the condition of subsurface structures such²⁶ as geological data and data from gravity method measurements. Gravity²⁷ method is one of the geophysical methods used to find out²⁸ conditions below the earth's²⁹ surface by measuring force field variations weight²⁷ of the earth.²⁹

The method used in hydrocarbon estimation was also revealed³⁰ by Sota (2011) in this research on estimating fault structures using the gravity method.

According to this research, the presence of³¹ faults can be associated³² with hydrocarbon traps. One of the methods used is the gravity method which was³³ chosen³³ for reasons of³⁴ response sensitivity, cheap³⁵ economically and technically in the field. The gravity method is very appropriate to use³⁶ for estimating fault locations because this method is able to³⁷ detect differences in density³⁸ contrast of rock bodies. Significant differences in rock density contrast indicate that the zone is a fault zone (Sota, 2011).

The gravity method is known to be able³⁹ to describe subsurface geological structures very well and has been widely used to identify faults,³⁹ and hydrocarbon prospect structures in oil and gas basins. Therefore, this method is very suitable for oil and gas exploration in the East Pondok Aren (EPN-001) field, Bekasi Regency, which is located⁴⁰ ⁴¹in the North West Java Basin.

2. LITERATURE REVIEW

2.1. Petroleum System

In oil and gas exploration activities in a basin, several elements are usually needed in a system ⁴² which is thought to contain hydrocarbons, ⁴² this system is usually called the Petroleum System. ⁴² Where the petroleum system includes several important elements, namely, source rock, reservoir rock, cap rock, trap and migration (Setyowiyoto, 2007; Permana, 2020).

2.1.1. Source Rock

There are three main source rocks in the North West Java Basin, ⁴³ namely lacustrine clay (oil-prone) ⁴³ which is reflected by the Banuwati Formation or Jatibarang Formation; deltaic fluvial coal and clay (oil and gas) reflected by the Upper Talang Akar Formation; and marine clay (bacterial gas) ⁴³ which is reflected by the Parigi Formation and Cisubuh Formation.

2.1.2. Reservoir Rock

The reservoir rocks in the Cibulakan Formation are primarily composed of sandstones, ⁴⁵ which have ⁴⁴ good porosity and permeability. These sandstones ⁴⁶ were deposited ⁴⁷ in a variety of environments, including fluvial, deltaic, and shallow marine settings. The Upper Cibulakan Formation often contains the best ⁴⁸ reservoir quality sands.

2.1.3. Cap Rock

The covering layer is an impermeable layer that can inhibit or cover the passage of hydrocarbons. The ⁴⁹ formation that acts as the main covering layer ⁴⁹ is the Cisubuh Formation, because this formation has impermeable lithology making ⁴⁹ it suitable as a barrier for hydrocarbons to migrate further.

2.1.4. Trap

The trap types in all petroleum systems in the⁵⁰ North West Java Basin are very similar.⁵¹ This is caused by the tectonic evolution of all sedimentary basins along the southern boundary of the Sunda Craton,⁵² so that the⁵³ types of geological structures and trapping mechanisms are almost the same. The main geological structures are wide⁵⁴ anticlinal domes and traps of tilted fault blocks. In⁵⁵ some areas with reservoir reef build-up, stratigraphic traps also play a role.⁵⁵

2.1.5. Migration

In the North West Java Basin, the main channels for lateral migration are mostly sandstone gaps that have a north-south orientation of the Talang Akar Formation, similar to the orientation of sandstone systems in the Main or Massive members (Upper Cibulakan Formation).⁵⁶ In this case, the fault becomes the main channel for vertical migration with fast transport (Prihantini, 2020).

2.2. Regional Geology

The North-West Java Basin is one of 128 sedimentary basins and is⁵⁷ known as a hydrocarbon basin. This basin is estimated to have reserves of 2.3 BBOE oil and 1.17 BBOE gas. The North West Java Basin is influenced⁵⁸ by a north-south trending block faulting system which⁵⁹ is divided⁶⁰ into grabens or several sub-basins from west to east, namely the Ciputat sub-basin, the Pasir Putih sub-basin and the Jatibarang sub-basin (Setiadi, 2018). The North West Java Basin has an age range from the Middle Eocene to the Quaternary. The regional startigraphic⁶¹ sequence from oldest to youngest is Bedrock, Jatibarang Formation, Lower Cibulakan Formation (Talang Akar, Baturaja),⁶² Upper Cibulakan Formation (Massive, Main, Pre -Parigi),^{61 63} Parigi Formation and⁶¹ Cisubuh Formation (Narpodo, 1996).

Sources of new oil and gas reserves in Tambelang District ⁶⁴are included in the Cibulakan Formation. The Cibulakan Formation consists of interbedded shale with sandstone and limestone. This formation ⁶⁵is divided into two members, namely Upper Cibulakan members and Lower Cibulakan members. The division of these members ⁶⁶is based on differences in depositional environments, where the Lower Cibulakan member is a transitional (paralic) deposit, ⁶⁷while the Upper Cibulakan member is a neritic deposit (Narpodo, 1996). Overall, the Cibulakan Formation is Early ⁶⁸Miocene to Middle Miocene in age (Rohmana, 2019).

⁶⁹If seen from the regional geological map (Figure 1), ⁶⁹the research area is included in floodplain deposits. ⁶⁹Generally, these floodplain deposits are dominated by alluvial deposits such as silt and mud, although fine sandstone occasionally appears which was deposited by stronger currents at the peak of the flood. The ⁷⁰rate of deposition is generally ⁷¹very low, ranging between 1 and 2 cm of silt-clay layer per flood period (Center for Water Resources Education and Training, 2017).

Figure 1. Geology of the Research Area (Ratman & Gafoer, 1998)

2.3. Gravity Method

The gravity method is a geophysical method that can describe subsurface geology based on variations in the ²⁹earth's gravitational field caused by density

differences between rocks (Pellokila, 2018) (Firdaus, 2018). This method has the advantage ⁷² that it can provide quite detailed information about the geological structure and density contrast of rocks (Kurniawan, 2022).

Knowledge of subsurface structures is ⁷³ important for planning exploration steps for oil and gas, geothermal ⁷⁴ and other minerals (Agussalim, 2019).

The basic principle of this gravitational method is to use Newton's law of gravity ⁷⁵ which states that the force of attraction between two points having masses m_1 and m_2 ⁷⁶ which ⁷⁷ are separated by a distance r , then the equation can be written:

$$F = G \frac{m_1 m_2}{r^2} \quad (1)$$

F is the force exerted between two particles with masses m_1 and m_2 , r is the distance between two particles, \hat{r} is the unit vector of m_1 and m_2 , and G is the universal gravitational constant ($6.6732 \times 10^{-11} \text{ Nm}^2/\text{kg}^2$) (Rizkiani, 2019).

2.4. Inversion Modeling

Inversion modeling is a field data processing process that involves mathematical and statistical solving techniques to obtain ⁷⁸ useful information regarding the distribution of subsurface physical properties. In the inversion process, field data ⁷⁹ is analyzed by performing curve fitting between the mathematical model and field data (Sihombing, 2018). In determining this inversion process, an initial model ⁸⁰ is needed in the form of average rock density (Maimuna, 2022).

⁸¹ The aim of the inversion process is to estimate previously unknown rock physical parameters. The inversion problem is formulated as an optimization problem where the object function of the model density ^{82 83} is minimized , according to boundary conditions within an error tolerance (Hiden, 2019).

Inversion modeling ^{84,85} is carried out by optimizing the singular value decomposition approach to gravity anomalies. This technique ⁸⁶ is based on the

linear inversion technique. The result of this inversion modeling is the distribution of rock density in two dimensions and is a picture of the subsurface of the research area (Setiadi, 2018).

3. METHODS

The research location is in the EPN-001 oil and gas drilling field in Tambelang District, Bekasi Regency, West Java with an area of around 1174.27 km² (Figure 2). The data used is gravity satellite data taken via the GGMPlus website with a total of 23,898 data consisting of gravity disturbance (gd), geoid, and topography.

GGMplus (Global Gravity Model plus) is a combination of GRACE and GOCE satellite gravity data, EGM2008 short-wavelength topographic gravity effects at a resolution of about 200 m for all terrestrial and nearshore regions of the Earth between $\pm 60^\circ$ latitude (Hirt, 2013). Gravitational disturbance data in GGMPlus is defined as the difference between Earth's gravity and normal gravity at the same point (Perozzi, 2021). And for the geoid it is the equipotential surface of the earth's real gravitational field that passes under the topographic mass and coincides with the mean sea level.

Figure 2. Research Location

The gravitational anomaly data obtained from the satellite is gravitational anomaly data that has been corrected using Free Air Correction (FAC), so the only corrections needed are terrain correction and Bouguer correction to obtain the Complete Bouguer Anomaly (CBA) value (Permana, 2022). Bouguer Correction is carried out to correct height by taking into account the effect of

the rock mass⁹⁷ around the measurement point. Terrain Correction is carried out⁹⁸ with the aim of bringing⁹⁹ the measured data to a flat plane so that the measured gravitational acceleration reading comes from the response of subsurface rocks (Pebrian, 2019). The final results¹⁰⁰ of data processing can describe 2D modeling of the earth's²⁹ subsurface at the EPN-001 oil and gas drill point¹¹ which is then correlated¹⁰² with the structural geological data of the research area (Figure 3).

Figure 3. Research Flow Chart

4. RESULTS & DISCUSSION

4.1. Topography Map

Figure 4. Topography Map

On the topographic map of the research area, it can be seen that the research¹⁰³ area has topographic values ranging from 0.02¹⁰³ – 27.65 m above sea level. The northern part of the research area has low topographic values ranging from 0.02¹⁰³ – 2.01 m because it is closer to the sea so the topographic value is very low,¹⁰³ in the southern part of the research area it has high topographic values ranging from 6.46¹⁰³ – 27.65 m. The topography of this study area is classified¹⁰⁴ as lowland morphology (Figure 4).

4.2. Complete Bouguer Anomaly Map

On the CBA map, the research area has gravity field anomaly values ranging from 36,860 – 53,342 mGal. The high anomaly has a value of 42,934 – 53,342 mGal, located in the northern and western parts of the study area which is^{105,107} thought¹⁰⁶ to be associated with volcanic rock deposits. The low anomaly has a value of 36,860 – 42,461 mGal, located in the south to east and a little in the

north ¹⁰⁸which ¹⁰⁹is thought to be associated with alluvium lithology and the Lower Cibulakan Formation ¹⁰⁸which is the target of research (black polygon) (Figure 5).

Figure 5. Complete Bouguer Anomaly Map

This CBA map still has ¹¹⁰quite high ambiguity because the anomaly that ¹¹¹is read is a combination of residual anomalies and regional anomalies, so this CBA map needs to be separated from the anomalies ¹¹²in order to eliminate this ambiguity.

4.3. Regional Anomaly Map

In the regional anomaly map the research area has anomaly values ranging from 36,817 – 53,623 mGal, it can be seen that the ¹¹³contour is smoother and ¹¹⁴has ¹¹⁵similarities to the CBA map. ¹¹⁶This is because regional anomalies describe ¹¹⁷very deep subsurface conditions, where the layers and rocks in the deep zone have homogeneous properties.

The high anomaly has values ranging from ¹¹⁷42,941 – 53,623 mGal in the northern and western parts of the study area. This high anomaly ¹¹⁸is thought to be associated with deep volcanic rock deposits. The low anomaly ¹¹⁹has values ranging from 36,817 – 42,179 mGal in the south to east and slightly in the north of the research area. This low anomaly ¹²⁰is thought to be associated with alluvium lithology and the Lower Cibulakan Formation in the inner zone ¹²¹which is the research target (Figure 6).

Figure 6. Regional Anomaly Map

4.4. Residual Anomaly Map

In the residual anomaly map, the research area has an anomaly value of -0.4463 – 0.3055 mGal. ¹²²It can be seen that the contour of the residual anomaly is more varied and slightly rougher ¹²²when compared to regional anomalies. ¹²³This is because residual anomalies describe shallow subsurface conditions or close

to the ground surface, where the layers and rocks in the shallow zone have heterogeneous properties.

The high anomaly has a value of 0.0056 - 0.3055 mGal ¹²⁴ which ¹²⁶ is located ¹²⁵ in the western, central ¹²⁴ and ^{124,126} eastern parts of the study area which ¹²⁷ extends from north to south. This high anomaly ¹²⁷ is thought to be associated with volcanic rock deposits. The low anomaly has a value of -0.4463 – (-0.0339) mGal ¹²⁸ which ¹³⁰ is located ¹²⁹ in almost all ¹³⁰ parts of the study area (Figure 7). This low anomaly ¹³¹ is thought to be associated with alluvium lithology and the Lower Cibulakan Formation, where in this ¹³² formation there is drilling point EPN-001 ¹³² which is the target of this research. Based on the residual anomaly map, a 2D model section ¹³³ will then be carried out.

Figure 7. Residual Anomaly Map

4.5. 2D Inversion Modeling

4.5.1. Section A-A'

In the 2D gravity model section A – A' which trends west to east with a depth of 4000 m with a root mean square error model value of 0.6, it can be seen that there are 3 lithologies or formations that make up the subsurface of the research area. The cap rock layer has a density contrast value of 1.5 – 1.8 g/cc at a depth of 0 – 2500 m ¹³⁴ which functions to resist the flow of oil and gas fluids to the surface. Layers ¹³⁵ consisting of volcanic rock have a density contrast value of 2.6 – 2.8 g/cc at a depth of 0 – 4000 m. The last layer, namely the Lower Cibulakan Formation, has a density contrast value of 2 – 2.3 g/cc at a depth of 2500 – 4000 m. This layer ¹³⁶ is thought to be ¹³⁷ a reservoir that stores hydrocarbons below the surface ¹³⁸ which is the target of this research (Figure 8).

Figure 8. 2D Gravity Inversion Model Section A-A'

4.5.2 Section B-B'

In the 2D gravity model section B – B', the drilling point cuts from west to east in the research area at a depth of 4000 m with a model root mean square error value of 0.39. The result is that there are ¹⁴⁰ 3 ¹³⁹ lithologies or formations that ¹⁴⁰ make up the subsurface of the research area. The cap rock layer has a density contrast value of 1.5 – 1.8 g/cc at a depth of ¹⁴¹ 0 – 2500 m. ¹⁴² Layers consisting of volcanic rock have a density contrast value of 2.6 – 2.8 g/cc at a depth of 0 – 4000 m. The last layer, namely the Lower Cibulakan Formation, has a density contrast value of 2 – 2.3 g/cc at a depth of 2500 – 4000 m. The Lower Cibulakan layer is thought to be ¹⁴³ a reservoir that stores hydrocarbons below the surface, where it can be seen ¹⁴⁴ that the drilling point that produces oil and gas fluids is at a depth of 2590 m (Figure 9).

Figure 9. 2D Gravity Inversion Model Section B - B'

4.5.3. Section C-C'

In the 2D gravity model section C – C' which ¹⁴⁵ trends west to east with a depth of 4000 m with a root mean square error model value of 1.19, it can be seen that there are ¹⁴⁶ 3 lithologies or formations that make up the subsurface of the research area. The cap rock layer has a density contrast value of 1.5 – 1.8 g/cc at a depth of 0 – 4000 m which functions ¹⁴⁷ to resist the flow of oil and gas fluids to the surface. Volcanic rock layers have a density contrast value of 2.6 – 2.8 g/cc at a depth of ¹⁴⁸ 0 – 4000 m. The last layer, namely the Lower Cibulakan Formation, has a density contrast value of 2.6 - 2.8 g/cc at a depth of 2500 –

4000 m. This layer ¹⁴⁹is thought to be ¹⁵⁰a reservoir that stores hydrocarbons below the surface ¹⁵¹which is the target of this research (Figure 10).

Figure 10. 2D Gravity Inversion Model Section C - C'

5. CONCLUSIONS

Based on the residual anomaly map, the low anomaly has a value of -0.4463 – (-0.0339) mGal ¹⁵²which is located in almost all parts of the study area. This low anomaly is thought to be associated with alluvium lithology and the Lower Cibulakan Formation, where ¹⁵²in this formation there is drilling point EPN-001 which is the target of this research.

In the three sections of the 2D model with a depth of 4000 m, ¹⁵⁴there are ¹⁵³3 lithologies or formations ¹⁵⁴that make up the subsurface of the research area. The cap rock layer has a density contrast value of 1.5 – 1.8 g/cc at a depth of 0 – 2500 m, the volcanic rock layer has a density contrast value of 2.6 – 2.8 g/cc at a depth of 0 – 4000 m, and ¹⁵⁵in the Lower Cibulakan Formation layer has density contrast value 2 – 2.3 g/cc at a depth of 2500 – 4000 m. The Lower Cibulakan Formation ¹⁵⁶is thought to be ¹⁵⁸a reservoir, where the drilling point at a depth of ^{157,158}2590 m and produces oil and gas fluids ^{157,158}which are visible in the 2D model section B - B'.

At the exploration stage, it is very important to collaborate with several geophysical methods to obtain more accurate and easier results at the data interpretation stage, especially the gravity method ¹⁵⁹to support seismic data in oil and gas exploration where it is quite difficult to see density parameters.

^{160,161} To increase oil and gas exploration, ^{161,162} various strategies can be implemented. The following is a potential strategy, namely Utilizing Geological and Geophysical Data by Building and maintaining a comprehensive ¹⁶⁰ database of ¹⁶³ geological and geophysical data ¹⁶³ to make it easier to identify potential areas.

REFERENCES

- Agussalim. (2019). Aplikasi Metode Gravitasi dengan Kontinuasi ke Atas (Upward Continuation) dalam Menginterpretasi Data Anomali Medan Gravitasi di Daerah Gunung Merapi. Jurnal Ilmu Fisika: Teori dan Aplikasinya, 1(1), 1-6.
- Dewi, ¹⁸ I.K., ¹⁶⁴ dkk. (2020). Pemodelan 3d Data Gravity Untuk Identifikasi Struktur Pembentukan Cekungan Hidrokarbon Wilayah Bajubang Provinsi Jambi. Jurnal Geofisika Eksplorasi, 6(3), 216-227. <https://doi.org/10.23960/jge.v6i3.10>.
- Direktorat Jenderal Minyak dan Gas Bumi. (2019). Statistik Minyak dan Gas Bumi 2019. Kementerian Energi dan Sumber Daya Mineral. <https://migas.esdm.go.id/cms/uploads/uploads/Statistik-Migas-2019---spread.pdf>.
- Direktorat Jenderal Minyak dan Gas Bumi. (2021). Statistik Minyak dan Gas Bumi Semester I 2021. Kementerian Energi dan Sumber Daya Mineral.
- Firdaus, S.A., Setianto, A. (2018). Interpretasi Struktur Geologi Berdasarkan Citra Landsat 8, Srtm Dan Anomali Medan Gravitasi Satelit Di Cekungan Jawa Timur Utara. Prosiding Seminar Nasional Geotik, 193-204.
- ¹⁶⁵ Hiden, dkk. (2019). Analisis dan Pemodelan Inversi Struktur Bawah Permukaan Berdasarkan Anomali Gravitasi Pulau Lombok. Indonesian Physical Review, 2(1), ¹⁶⁶ 1-8.
- Hirt, C, ¹⁸ S.J. Claessens, T. Fecher, M. Kuhn, R. Pail, M. Rexer ¹⁶⁷ (2013), New ultra-high resolution picture of ²⁹ Earth's gravity field. Geophysical Research Letters, 40. Doi: 10.1002/grl.50838.

Ira, L. (2023). Cerita Pertamina Temukan Sumber Minyak Baru di Bekasi, Pernah Terjadi di Lokasi Lain. Diakses pada 1 Maret 2024, <https://bisnis.tempo.co/read/1812576/cerita-pertamina-temukan-sumber-minyak-baru-di-bekasipernah-terjadi-di-lokasi-lain>.

Kurniawan, dkk.¹⁶⁸ (2022). Identifikasi Kemenerusan Sesar Dan Potensi Reservoir Air Panas Cagar Dengan Menggunakan Metode Gravitasi. Jurnal Pendidikan Fisika dan Terapannya, 5(1), 21-27.

Maimuna, A.K., Miftakhunnis, A.F., Segoro, Y.A. (2022). Pemodelan Inversi 3 Dimensi Untuk Identifikasi Dugaan Keberadaan Sesar Menggunakan Data Anomali Gaya Berat Di Laut Flores (Studi Kasus Gempa Flores 14 Desember 2021). Progress: Jurnal Geofisika, 1(1), 54-61.

Narpodo, J. (1996). Studi Konversi Kedalaman dengan Metode Stacking Velocity dan Layer Cake di daerah Jawa Barat Utara. Skripsi-S1 Geofisika FMIPA UGM, Yogyakarta.

Pebrian, M. Q., Marwan, Ismail. (2019). Interpretasi Data Anomali Gravitasi untuk Identifikasi Endapan Aluvium di Kuala Gingjeng, Aceh Besar. Risalah Fisika, 3(1), 15-19.

Pellokila, A.I., Bernandus, Tanesib, J.L. (2018). Identifikasi Keberadaan Basement Di Bawah Cekungan Timor Berdasarkan Data Anomali Gravitasi Dengan Pemodelan Tiga Dimensi. Jurnal Fisika, 3(1), 1-11.

Permana, N.P., Gunawan, B., Nafian, M. (2022)¹⁶⁹, Analysis of Subsurface Structure of Sembalun Geothermal Prospect Area, East Lombok with 2D and 3D Gravity Modeling. Al-Fiziya: Journal of Materials Science, Geophysics, Instrumentation and Theoretical Physics, 5(1), 10 – 22.

Permana, N.R. (2020). Pemetaan Struktur Batuan Dasar Menggunakan Metode Energy Spectral Analysis – Multi Window Test (ESA – MWT) Data Gravitasi Pada

Sub Cekungan Sakala. Program Studi Fisika, Fakultas Sains dan Teknologi, UIN Syarif Hidayatullah Jakarta.

Perozzi, L., Guglielmetti, L., Moscariello, A. (2021). Quantitative uncertainty analysis of gravity disturbance. The case of the Geneva Basin (Switzerland). *Journal of Applied Geophysics*, 193. Doi: 10.1016/j.jappgeo.2021.104431.

Prasetiyohadi, S.A., dkk. (2022). Evaluasi Dan Korelasi Batuan Induk-Minyak Bumi Berdasarkan Analisis Geokimia Hidrokarbon Di Lapangan Sp, Sub-Cekungan Jambi. *Padjadjaran Geoscience Journal*, 6(3), 855-875.

Purnama, D.I., ¹⁷⁰ dkk. (2020). Identifikasi Potensi Batuan Induk pada Formasi Santul di Sub-Cekungan Tarakan, Kalimantan Utara. *Prisma Fisika*, 8(1), 1-4.

Pusat Pendidikan dan Pelatihan Sumber Daya Air dan Konstruksi. (2017). Modul Pengelolaan Dataran Banjir Pelatihan Pengendalian Banjir. Kementrian Pekerjaan Umum dan Perumahan Rakyat.

Rahmayanti, L., Rahmah, D.M., Larashati. (2021). Analisis Pemanfaatan Sumber Daya Energi Minyak Dan Gas Bumi Di Indonesia. *Jurnal Sains Edukatika Indonesia (JSEI)*, 3(2), 9-16.

Ratman, N., Gafoer, S. (1998). Peta Geologi Lembar Jawa Bagian Barat. Pusat Penelitian dan pengembangan Geologi.

Rizkiani, D.N, Rustadi. (2019). Interpretasi Sistem Panas Bumi Suwawa Berdasarkan Data Gaya Berat. *Jurnal Geofisika Eksplorasi*, 5(2), 44-54.

Rohmana, S.C., ¹⁷¹ dkk. (2019). Identifikasi Reservoar Hidrokarbon Resistivitas Rendah: Studi Kasus Di Formasi Cibulakan Atas, Cekungan Jawa Barat Utara Dan Formasi Gumai, Sub-Cekungan Jambi. *Prosiding TAU SNAR-TEK 2019 Seminar Nasional Rekayasa dan Teknologi*, 178-184.

Rosid, M.S., Tullailah, N.N., Wibowo, R.A. (2020). Identifikasi potensi jebakan hidrokarbon di daerah "X" sedimen pra-tercier cekungan Sumatra Utara menggunakan data gravitasi dan seismik. *Jurnal Fisika*, 10(1), 8-21.

Setiadi, I., Pratama., A.W. (2018). Pola Struktur dan Konfigurasi Geologi Bawah Permukaan Cekungan Jawa Barat Utara Berdasarkan Analisis Gayaberat
Structural Pattern and Subsurface Geological Configuration of North West Java Basin Based on Gravity Analysis. Jurnal Geologi dan Sumberdaya Mineral, 19(2), 59 – 72.

Setyowiyoto, J., Datun, M., Winardi, S. (2007). Geologi Dan Tinjauan Petroleum System Daerah Bancak Kabupaten Semarang Berdasarkan Manifestasi Permukaan. Media Teknik, No. 1.

Sihombing. R.B., Sarkowi, H.M., Rustadi. (2018). Pemodelan Dan Analisa Struktur Bawah Permukaan Daerah Prospek Panasbumi Kepahiang Berdasarkan Metode Gayaberat. Jurnal Geofisika Eksplorasi, 4(2).

Sota, I. (2011). Pendugaan Struktur Patahan Dengan Metode Gayaberat. POSITRON, 1(1), 25-30.

Tim CNN Indonesia. (2023). Pertamina Temukan Sumur Migas Baru di Bekasi. CNN Indonesia. (Diakses pada 3 Maret 2024), <https://www.cnnindonesia.com/ekonomi/20231219112029-85-1039096/pertamina-temukan-sumur-migas-baru-di-bekasi>.

1.	which was	Wordy sentences	Clarity
2.	needs to → must	Wordy sentences	Clarity
3.	<i>The drill data needs to be correlated with other supporting data such as geophysical methods to ensure the accuracy.</i>	Ungrammatical sentence	Correctness
4.	<i>was conducted</i>	Passive voice misuse	Clarity
5.	examined	Wordy sentences	Clarity
6.	<i>was taken</i>	Passive voice misuse	Clarity
7.	bouguer → Bouguer	Misspelled words	Correctness
8.	<i>were carried out</i>	Passive voice misuse	Clarity
9.	which	Punctuation in compound/complex sentences	Correctness
10.	<i>are thought</i>	Passive voice misuse	Clarity
11.	<i>be seen</i>	Passive voice misuse	Clarity
12.	<i>As a result, continuous oil production and oil reserves will increasingly run out, causing the oil exploration process to stop, resulting in old wells (Rosid, 2020).</i>	Incorrect phrasing	Correctness
13.	important → essential, vital, crucial	Word choice	Engagement
14.	in an effort to → to	Wordy sentences	Clarity
15.	<i>been operated</i>	Passive voice misuse	Clarity
16.	<i>been exploited</i>	Passive voice misuse	Clarity
17.	also	Wordy sentences	Clarity

18.	<i>PT; I.K.; S.J.</i>	Text inconsistencies	Correctness
19.	<i>which</i>	Punctuation in compound/complex sentences	Correctness
20.	<i>,and is → it is</i>	Hard-to-read text	Clarity
21.	<i>is included</i>	Passive voice misuse	Clarity
22.	<i>18</i>	Comma misuse within clauses	Correctness
23.	<i>1,</i>	Punctuation in compound/complex sentences	Correctness
24.	<i>To utilize the energy potential of oil and gas</i>	Misplaced words or phrases	Correctness
25.	<i>been obtained</i>	Passive voice misuse	Clarity
26.	<i>such</i>	Punctuation in compound/complex sentences	Correctness
27.	<i>Gravity method is one of the geophysical methods used to find out conditions below the earth's surface by measuring force field variations weight of the earth.</i>	Ungrammatical sentence	Correctness
28.	<i>out</i>	Wordy sentences	Clarity
29.	<i>earth's; earth; Earth; Earth's</i>	Text inconsistencies	Correctness
30.	<i>was also revealed</i>	Passive voice misuse	Clarity
31.	<i>the presence of</i>	Wordy sentences	Clarity
32.	<i>be associated</i>	Passive voice misuse	Clarity
33.	<i>was chosen</i>	Passive voice misuse	Clarity
34.	<i>reasons of</i>	Wordy sentences	Clarity

35.	<i>One of the methods used is the gravity method which was chosen for reasons of response sensitivity, cheap economically and technically in the field.</i>	Incorrect phrasing	Correctness
36.	to use	Wordy sentences	Clarity
37.	is able to → can	Wordy sentences	Clarity
38.	the density	Determiner use (a/an/the/this, etc.)	Correctness
39.	<i>The gravity method is known to be able to describe subsurface geological structures very well and has been widely used to identify faults, and hydrocarbon prospect structures in oil and gas basins.</i>	Paragraph can be perfected	Clarity
40.	which is	Wordy sentences	Clarity
41.	<i>is located</i>	Passive voice misuse	Clarity
42.	<i>In oil and gas exploration activities in a basin, several elements are usually needed in a system which is thought to contain hydrocarbons, this system is usually called the Petroleum System. Where the petroleum system includes several important elements, namely, source rock, reservoir rock, cap ro...</i>	Paragraph can be perfected	Clarity
43.	<i>There are three main source rocks in the North West Java Basin, namely lacustrine clay (oil-prone) which is reflected by the Banuwati Formation or Jatibarang Formation; deltaic fluvial coal and clay (oil and gas) reflected by the Upper Talang Akar Formation; and marine clay (bacterial gas) which is...</i>	Paragraph can be perfected	Clarity
44.	good → suitable	Word choice	Engagement

45.	<i>The reservoir rocks in the Cibulakan Formation are primarily composed of sandstones, which have good porosity and permeability.</i>	Unclear sentences	Clarity
46.	<i>were deposited</i>	Passive voice misuse	Clarity
47.	a variety of → various	Wordy sentences	Clarity
48.	reservoir-quality	Misspelled words	Correctness
49.	<i>The formation that acts as the main covering layer is the Cisubuh Formation, because this formation has impermeable lithology making it suitable as a barrier for hydrocarbons to migrate further.</i>	Paragraph can be perfected	Clarity
50.	<i>The trap types in all petroleum systems in the North West Java Basin are very similar.</i>	Unclear sentences	Clarity
51.	<i>This</i>	Intricate text	Clarity
52.	Craton,	Punctuation in compound/complex sentences	Correctness
53.	that the	Wordy sentences	Clarity
54.	wide → vast	Word choice	Engagement
55.	<i>In some areas with reservoir reef build-up, stratigraphic traps also play a role.</i>	Unclear sentences	Clarity
56.		Incorrect citation format	Correctness
57.	and is	Wordy sentences	Clarity
58.	<i>is influenced</i>	Passive voice misuse	Clarity
59.	which	Punctuation in compound/complex sentences	Correctness

60.	<i>is divided</i>	Passive voice misuse	Clarity
61.	<i>The regional stratigraphic sequence from oldest to youngest is Bedrock, Jatibarang Formation, Lower Cibulakan Formation (Talang Akar, Baturaja), Upper Cibulakan Formation (Massive, Main, Pre -Parigi), Parigi Formation and Cisubuh Formation (Narpodo, 1996).</i>	Ungrammatical sentence	Correctness
62.		Incorrect citation format	Correctness
63.		Incorrect citation format	Correctness
64.	<i>are included</i>	Passive voice misuse	Clarity
65.	<i>is divided</i>	Passive voice misuse	Clarity
66.	<i>is based</i>	Passive voice misuse	Clarity
67.	,while → In contrast	Hard-to-read text	Clarity
68.	Miocene	Wordy sentences	Clarity
69.	<i>If seen from the regional geological map (Figure 1), the research area is included in floodplain deposits. Generally, these floodplain deposits are dominated by alluvial deposits such as silt and mud, although fine sandstone occasionally appears which was deposited by stronger currents at the peak ...</i>	Paragraph can be perfected	Clarity
70.	deposition rate	Wordy sentences	Clarity
71.	very low → meager, shallow	Word choice	Engagement

72.	<i>This method has the advantage that it can provide quite detailed information about the geological structure and density contrast of rocks (Kurniawan, 2022).</i>	Unclear sentences	Clarity
73.	important → essential, vital	Word choice	Engagement
74.	, and	Comma misuse within clauses	Correctness
75.	which	Punctuation in compound/complex sentences	Correctness
76.	which	Punctuation in compound/complex sentences	Correctness
77.	are separated	Passive voice misuse	Clarity
78.	useful → helpful	Word choice	Engagement
79.	is analyzed	Passive voice misuse	Clarity
80.	is needed	Passive voice misuse	Clarity
81.	The inversion process aims	Wordy sentences	Clarity
82.	is minimized	Passive voice misuse	Clarity
83.	minimized,	Punctuation in compound/complex sentences	Correctness
84.	is carried out	Passive voice misuse	Clarity
85.	carried out → done	Wordy sentences	Clarity
86.	is based	Passive voice misuse	Clarity
87.	with	Punctuation in compound/complex sentences	Correctness
88.	a total of	Wordy sentences	Clarity




89.	<i>The data used is gravity satellite data taken via the GGMPPlus website with a total of 23,898 data consisting of gravity disturbance (gd), geoid, and topography.</i>	Incorrect phrasing	Correctness
90.	<i>is defined</i>	Passive voice misuse	Clarity
91.	And → Moreover, Furthermore,	Inappropriate colloquialisms	Delivery
92.	<i>And for the geoid it is the equipotential surface of the earth's real gravitational field that passes under the topographic mass and coincides with the mean sea level.</i>	Incorrect phrasing	Correctness
93.	real → actual, natural	Word choice	Engagement
94.	<i>And for the geoid it is the equipotential surface of the earth's real gravitational field that passes under the topographic mass and coincides with the mean sea level.</i>	Unclear sentences	Clarity
95.	<i>been corrected</i>	Passive voice misuse	Clarity
96.	<i>is carried out</i>	Passive voice misuse	Clarity
97.	<i>Bouguer Correction is carried out to correct height by taking into account the effect of the rock mass around the measurement point.</i>	Unclear sentences	Clarity
98.	<i>is carried out</i>	Passive voice misuse	Clarity
99.	to bring	Wordy sentences	Clarity
100.	data processing results	Wordy sentences	Clarity
101.	which	Punctuation in compound/complex sentences	Correctness

102.	<i>is then correlated</i>	Passive voice misuse	Clarity
103.	<i>On the topographic map of the research area, it can be seen that the research area has topographic values ranging from 0.02 – 27.65 m above sea level. The northern part of the research area has low topographic values ranging from 0.02 - 2.01 m because it is closer to the sea so the topographic valu...</i>	Paragraph can be perfected	Clarity
104.	<i>is classified</i>	Passive voice misuse	Clarity
105.	<i>which</i>	Punctuation in compound/complex sentences	Correctness
106.	<i>is thought</i>	Passive voice misuse	Clarity
107.	<i>The high anomaly has a value of 42,934 - 53,342 mGal, located in the northern and western parts of the study area which is thought to be associated with volcanic rock deposits.</i>	Unclear sentences	Clarity
108.	<i>The low anomaly has a value of 36,860 – 42,461 mGal, located in the south to east and a little in the north which is thought to be associated with alluvium lithology and the Lower Cibulakan Formation which is the target of research (black polygon) (Figure 5).</i>	Ungrammatical sentence	Correctness
109.	<i>is thought</i>	Passive voice misuse	Clarity
110.	<i>quite</i> → <i>relatively</i>	Word choice	Engagement
111.	<i>is read</i>	Passive voice misuse	Clarity
112.	<i>in order to</i> → <i>to</i>	Wordy sentences	Clarity

113.	<i>In the regional anomaly map the research area has anomaly values ranging from 36,817 – 53,623 mGal, it can be seen that the contour is smoother and has similarities to the CBA map.</i>	Paragraph can be perfected	Clarity
114.	<i>This</i>	Intricate text	Clarity
115.	This is because regional	Wordy sentences	Clarity
116.	very deep → intense	Word choice	Engagement
117.	42,941 → 42,941 to	Wrong or missing prepositions	Correctness
118.	<i>is thought</i>	Passive voice misuse	Clarity
119.	has values ranging → ranges	Wordy sentences	Clarity
120.	<i>is thought</i>	Passive voice misuse	Clarity
121.	which	Punctuation in compound/complex sentences	Correctness
122.	<i>It can be seen that the contour of the residual anomaly is more varied and slightly rougher when compared to regional anomalies.</i>	Paragraph can be perfected	Clarity
123.	<i>This</i>	Intricate text	Clarity
124.	<i>The high anomaly has a value of 0.0056 - 0.3055 mGal which is located in the western, central and eastern parts of the study area which extends from north to south.</i>	Ungrammatical sentence	Correctness
125.	<i>is located</i>	Passive voice misuse	Clarity

126.	<i>The high anomaly has a value of 0.0056 - 0.3055 mGal which is located in the western, central and eastern parts of the study area which extends from north to south.</i>	Unclear sentences	Clarity
127.	<i>is thought</i>	Passive voice misuse	Clarity
128.	<i>which</i>	Punctuation in compound/complex sentences	Correctness
129.	<i>is located</i>	Passive voice misuse	Clarity
130.	<i>The low anomaly has a value of -0.4463 – (-0.0339) mGal which is located in almost all parts of the study area (Figure 7).</i>	Unclear sentences	Clarity
131.	<i>is thought</i>	Passive voice misuse	Clarity
132.	<i>This low anomaly is thought to be associated with alluvium lithology and the Lower Cibulakan Formation, where in this formation there is drilling point EPN-001 which is the target of this research.</i>	Ungrammatical sentence	Correctness
133.	<i>be carried out</i>	Passive voice misuse	Clarity
134.	<i>The cap rock layer has a density contrast value of 1.5 – 1.8 g/cc at a depth of 0 – 2500 m which functions to resist the flow of oil and gas fluids to the surface.</i>	Incorrect phrasing	Correctness
135.	consisting	Wordy sentences	Clarity
136.	<i>is thought</i>	Passive voice misuse	Clarity
137.	thought to be → <i>considered</i>	Wordy sentences	Clarity

138.	which	Punctuation in compound/complex sentences	Correctness
139.	3 → three	Improper formatting	Correctness
140.	<i>The result is that there are 3 lithologies or formations that make up the subsurface of the research area.</i>	Unclear sentences	Clarity
141.	<i>The cap rock layer has a density contrast value of 1.5 – 1.8 g/cc at a depth of 0 – 2500 m.</i>	Unclear sentences	Clarity
142.	consisting	Wordy sentences	Clarity
143.	<i>is thought</i>	Passive voice misuse	Clarity
144.	<i>be seen</i>	Passive voice misuse	Clarity
145.	which	Punctuation in compound/complex sentences	Correctness
146.	3 → three	Improper formatting	Correctness
147.	<i>The cap rock layer has a density contrast value of 1.5 – 1.8 g/cc at a depth of 0 – 4000 m which functions to resist the flow of oil and gas fluids to the surface.</i>	Incorrect phrasing	Correctness
148.	<i>Volcanic rock layers have a density contrast value of 2.6 – 2.8 g/cc at a depth of 0 – 4000 m.</i>	Unclear sentences	Clarity
149.	<i>is thought</i>	Passive voice misuse	Clarity
150.	thought to be → considered	Wordy sentences	Clarity
151.	which	Punctuation in compound/complex sentences	Correctness

152.	<i>Based on the residual anomaly map, the low anomaly has a value of -0.4463 – (-0.0339) mGal which is located in almost all parts of the study area. This low anomaly is thought to be associated with alluvium lithology and the Lower Cibulakan Formation, where in this formation there is drilling point ...</i>	Paragraph can be perfected	Clarity
153.	 → 	Improper formatting	Correctness
154.	<i>In the three sections of the 2D model with a depth of 4000 m, there are 3 lithologies or formations that make up the subsurface of the research area.</i>	Unclear sentences	Clarity
155.		Wrong or missing prepositions	Correctness
156.	<i>is thought</i>	Passive voice misuse	Clarity
157.	<i>The Lower Cibulakan Formation is thought to be a reservoir, where the drilling point at a depth of 2590 m and produces oil and gas fluids which are visible in the 2D model section B - B'.</i>	Ungrammatical sentence	Correctness
158.	<i>The Lower Cibulakan Formation is thought to be a reservoir, where the drilling point at a depth of 2590 m and produces oil and gas fluids which are visible in the 2D model section B - B'.</i>	Unclear sentences	Clarity
159.	<i>At the exploration stage, it is very important to collaborate with several geophysical methods to obtain more accurate and easier results at the data interpretation stage, especially the gravity method to support seismic data in oil and gas exploration where it is quite difficult to see density par...</i>	Paragraph can be perfected	Clarity

160.	<i>To increase oil and gas exploration, various strategies can be implemented. The following is a potential strategy, namely Utilizing Geological and Geophysical Data by Building and maintaining a comprehensive database of geological and geophysical data to make it easier to identify potential areas.</i>	Paragraph can be perfected	Clarity
161.	<i>To increase oil and gas exploration, various strategies can be implemented.</i>	Incorrect phrasing	Correctness
162.	<i>be implemented</i>	Passive voice misuse	Clarity
163.	<i>The following is a potential strategy, namely Utilizing Geological and Geophysical Data by Building and maintaining a comprehensive database of geological and geophysical data to make it easier to identify potential areas.</i>	Unclear sentences	Clarity
164.	<i>dkk</i>	Unknown words	Correctness
165.		Incorrect citation format	Correctness
166.		Incorrect citation format	Correctness
167.		Incorrect citation format	Correctness
168.		Incorrect citation format	Correctness
169.		Incorrect citation format	Correctness
170.	<i>dkk</i>	Unknown words	Correctness
171.	<i>dkk</i>	Unknown words	Correctness